

LOW DIMENSIONAL d -ALGEBRAS

AARON KAUFER

MENTOR: LUCAS MASON-BROWN

MAY 20 2017

PRIMES CONFERENCE

What is an Algebra?

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- 2 Multiplication: If $a, b \in A$, then $a \cdot b \in A$.

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- 2 Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- 3 Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- 2 Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- 3 Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- 2 Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- 3 Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

- \mathbb{Q} : The rational numbers

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- 1 Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- 2 Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- 3 Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

- \mathbb{Q} : The rational numbers
- \mathbb{R} : The real numbers

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

- \mathbb{Q} : The rational numbers
- \mathbb{R} : The real numbers
- \mathbb{C} : The complex numbers

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

- \mathbb{Q} : The rational numbers
- \mathbb{R} : The real numbers
- \mathbb{C} : The complex numbers
- $\mathbb{Z}/p\mathbb{Z}$: The integers modulo a prime.

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

Here, a scalar means an element of a field F . F could be:

- \mathbb{Q} : The rational numbers
- \mathbb{R} : The real numbers
- \mathbb{C} : The complex numbers
- $\mathbb{Z}/p\mathbb{Z}$: The integers modulo a prime.

If F is the field of scalars for an algebra A , then we say that A is an algebra over F .

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

$$\bullet 0 \in A \text{ and } 1 \in A$$

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $1 \cdot a = a$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $1 \cdot a = a$
- $a + (-a) = 0$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $1 \cdot a = a$
- $a + (-a) = 0$
- $(a + b) \cdot c = a \cdot c + b \cdot c$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $1 \cdot a = a$
- $a + (-a) = 0$
- $(a + b) \cdot c = a \cdot c + b \cdot c$
- $a \cdot (b + c) = a \cdot b + a \cdot c$

What is an Algebra?

Definition

An **Algebra** is a set A together with three operations:

- ➊ Addition/Subtraction: If $a, b \in A$, then $a + b \in A$ and $a - b \in A$.
- ➋ Multiplication: If $a, b \in A$, then $a \cdot b \in A$.
- ➌ Scalar multiplication: If $a \in A$ and k is a scalar, then $ka \in A$.

For $a, b, c \in A$ and k is a scalar, we must have:

- $a + b = b + a$
- $(a + b) + c = a + (b + c)$
- $(a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $0 \in A$ and $1 \in A$
- $0 + a = a$
- $1 \cdot a = a$
- $a + (-a) = 0$
- $(a + b) \cdot c = a \cdot c + b \cdot c$
- $a \cdot (b + c) = a \cdot b + a \cdot c$
- $k(a \cdot b) = (ka) \cdot b = a \cdot (kb)$

Examples of Algebras

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$
 - ▶ Scalar Multiplication: $7(1 + 2i) = 7 + 14i$

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$
 - ▶ Scalar Multiplication: $7(1 + 2i) = 7 + 14i$
- Polynomials over a field form an algebra. For example:

$$\mathbb{Q}[x] = \{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \mid a_0, \dots, a_n \in \mathbb{Q}\}$$

is an algebra with field of scalars \mathbb{Q} .

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$
 - ▶ Scalar Multiplication: $7(1 + 2i) = 7 + 14i$
- Polynomials over a field form an algebra. For example:

$$\mathbb{Q}[x] = \{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \mid a_0, \dots, a_n \in \mathbb{Q}\}$$

is an algebra with field of scalars \mathbb{Q} .

- ▶ Addition: $(3x^2 + 1) + (2x + 5) = 3x^2 + 2x + 6$

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$
 - ▶ Scalar Multiplication: $7(1 + 2i) = 7 + 14i$
- Polynomials over a field form an algebra. For example:

$$\mathbb{Q}[x] = \{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \mid a_0, \dots, a_n \in \mathbb{Q}\}$$

is an algebra with field of scalars \mathbb{Q} .

- ▶ Addition: $(3x^2 + 1) + (2x + 5) = 3x^2 + 2x + 6$
- ▶ Multiplication: $(3x^2 + 1) \cdot (2x + 5) = 6x^3 + 15x^2 + 2x + 5$

Examples of Algebras

- The complex numbers \mathbb{C} is an algebra with field of scalars \mathbb{R} .
 - ▶ Addition: $(1 + 2i) + (2 + 6i) = 3 + 9i$
 - ▶ Multiplication: $(1 + 2i) \cdot (2 + 6i) = -10 + 10i$
 - ▶ Scalar Multiplication: $7(1 + 2i) = 7 + 14i$
- Polynomials over a field form an algebra. For example:

$$\mathbb{Q}[x] = \{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \mid a_0, \dots, a_n \in \mathbb{Q}\}$$

is an algebra with field of scalars \mathbb{Q} .

- ▶ Addition: $(3x^2 + 1) + (2x + 5) = 3x^2 + 2x + 6$
- ▶ Multiplication: $(3x^2 + 1) \cdot (2x + 5) = 6x^3 + 15x^2 + 2x + 5$
- ▶ Scalar multiplication: $\frac{1}{3}(2x + 5) = \frac{2}{3}x + \frac{5}{3}$

Commutative Algebras

- All three algebras from the previous slide had the property that multiplication was **commutative**. This means that the equation:

$$a \cdot b = b \cdot a \text{ for all } a, b \in A$$

is true when $A = \mathbb{C}$ or $A = \mathbb{Q}[x]$ (or any other polynomial algebra).

- All three algebras from the previous slide had the property that multiplication was **commutative**. This means that the equation:

$$a \cdot b = b \cdot a \text{ for all } a, b \in A$$

is true when $A = \mathbb{C}$ or $A = \mathbb{Q}[x]$ (or any other polynomial algebra).

Definition

An algebra A is called a **Commutative Algebra** if:

$$a \cdot b = b \cdot a$$

for all $a, b \in A$.

Noncommutative Algebras

Not all algebras are commutative!

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

► Addition: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 7 & 7 \end{pmatrix}$

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

- ▶ Addition: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 7 & 7 \end{pmatrix}$
- ▶ Multiplication: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} \cdot \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 59 & 36 \\ 13 & 9 \end{pmatrix}$

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

- ▶ Addition: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 7 & 7 \end{pmatrix}$
- ▶ Multiplication: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} \cdot \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 59 & 36 \\ 13 & 9 \end{pmatrix}$
- ▶ Scalar Multiplication: $6 \begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 6 & 42 \\ -6 & 12 \end{pmatrix}$

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

- ▶ Addition: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 7 & 7 \end{pmatrix}$
 - ▶ Multiplication: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} \cdot \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 59 & 36 \\ 13 & 9 \end{pmatrix}$
 - ▶ Scalar Multiplication: $6 \begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 6 & 42 \\ -6 & 12 \end{pmatrix}$
- ▶ To see that $M_{2,2}(\mathbb{R})$ is not commutative, set $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$.

Noncommutative Algebras

Not all algebras are commutative!

- Consider:

$$M_{2,2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$$

We have our familiar operations:

► Addition: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} + \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 4 & 8 \\ 7 & 7 \end{pmatrix}$

► Multiplication: $\begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} \cdot \begin{pmatrix} 3 & 1 \\ 8 & 5 \end{pmatrix} = \begin{pmatrix} 59 & 36 \\ 13 & 9 \end{pmatrix}$

► Scalar Multiplication: $6 \begin{pmatrix} 1 & 7 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 6 & 42 \\ -6 & 12 \end{pmatrix}$

- To see that $M_{2,2}(\mathbb{R})$ is not commutative, set $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$. Then:

$$A \cdot B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \text{ but } B \cdot A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

Dimension

Dimension

Suppose A is an algebra over F .

Dimension

Suppose A is an algebra over F .

Definition

Suppose a_1, \dots, a_n are all elements of A . A **linear combination** of a_1, \dots, a_n is a finite sum of the form:

$$k_1 a_1 + k_2 a_2 + \cdots + k_n a_n$$

where k_1, \dots, k_n are all scalars.

Dimension

Suppose A is an algebra over F .

Definition

Suppose a_1, \dots, a_n are all elements of A . A **linear combination** of a_1, \dots, a_n is a finite sum of the form:

$$k_1 a_1 + k_2 a_2 + \cdots + k_n a_n$$

where k_1, \dots, k_n are all scalars.

For example, set $A = M_{2,2}(\mathbb{R})$ and $F = \mathbb{R}$. Then the matrix $\begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix}$ is a linear combination of the matrices $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ because:

$$\begin{pmatrix} 5 & 0 \\ 0 & 2 \end{pmatrix} = 5 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + 2 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

Dimension

- As a matter of fact, every matrix in $M_{2,2}(\mathbb{R})$ is a linear combination of the matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

because:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + d \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

- As a matter of fact, every matrix in $M_{2,2}(\mathbb{R})$ is a linear combination of the matrices:

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

because:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = a \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + b \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} + c \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} + d \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

- Likewise, every element of the algebra \mathbb{C} is a linear combination of the numbers 1 and i , because:

$$a + bi = a(1) + b(i)$$

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

For example:

- 1 and i span \mathbb{C} .

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

For example:

- 1 and i span \mathbb{C} .
- $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$ and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ span $M_{2,2}(\mathbb{R})$

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

For example:

- 1 and i span \mathbb{C} .
- $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ span $M_{2,2}(\mathbb{R})$

Definition

The **dimension** of A is the minimum number of elements in A needed to span A . The dimension of A is denoted $\dim(A)$.

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

For example:

- 1 and i span \mathbb{C} .
- $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ span $M_{2,2}(\mathbb{R})$

Definition

The **dimension** of A is the minimum number of elements in A needed to span A . The dimension of A is denoted $\dim(A)$.

For example:

- $\dim(\mathbb{C}) = 2$

Dimension

Suppose A is an algebra over F .

Definition

A set of elements $a_1, \dots, a_n \in A$ is said to **span** A if every element of A can be written as a linear combination of a_1, \dots, a_n .

For example:

- 1 and i span \mathbb{C} .
- $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$ and $\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ span $M_{2,2}(\mathbb{R})$

Definition

The **dimension** of A is the minimum number of elements in A needed to span A . The dimension of A is denoted $\dim(A)$.

For example:

- $\dim(\mathbb{C}) = 2$
- $\dim(M_{2,2}(\mathbb{R})) = 4$

What is a d -Algebra?

What is a d -Algebra?

Definition

A d -Algebra over a field F is a regular algebra A together with a function d , called the “differential” of A , which satisfies the properties:

What is a d -Algebra?

Definition

A d -Algebra over a field F is a regular algebra A together with a function d , called the “differential” of A , which satisfies the properties:

- 1 d is linear:

$$d(a + b) = d(a) + d(b) \text{ and } d(ka) = kd(a)$$

whenever $a, b \in A$ and $k \in F$.

What is a d -Algebra?

Definition

A d -Algebra over a field F is a regular algebra A together with a function d , called the “differential” of A , which satisfies the properties:

- 1 d is linear:

$$d(a + b) = d(a) + d(b) \text{ and } d(ka) = kd(a)$$

whenever $a, b \in A$ and $k \in F$.

- 2 d is a derivation:

$$d(a \cdot b) = d(a) \cdot b + a \cdot d(b)$$

for all $a, b \in A$.

What is a d -Algebra?

Definition

A d -Algebra over a field F is a regular algebra A together with a function d , called the “differential” of A , which satisfies the properties:

- ① d is linear:

$$d(a + b) = d(a) + d(b) \text{ and } d(ka) = kd(a)$$

whenever $a, b \in A$ and $k \in F$.

- ② d is a derivation:

$$d(a \cdot b) = d(a) \cdot b + a \cdot d(b)$$

for all $a, b \in A$.

- ③ $d^2 = 0$:

$$d(d(a)) = 0 \text{ for all } a \in A$$

What is a d -Algebra?

Definition

A d -Algebra over a field F is a regular algebra A together with a function d , called the “differential” of A , which satisfies the properties:

- ① d is linear:

$$d(a + b) = d(a) + d(b) \text{ and } d(ka) = kd(a)$$

whenever $a, b \in A$ and $k \in F$.

- ② d is a derivation:

$$d(a \cdot b) = d(a) \cdot b + a \cdot d(b)$$

for all $a, b \in A$.

- ③ $d^2 = 0$:

$$d(d(a)) = 0 \text{ for all } a \in A$$

- ④ d -commutativity:

$$a \cdot b = b \cdot a + d(b) \cdot d(a)$$

Examples of d -algebras

- Trivial d -algebra:

- Trivial d -algebra:
 - ▶ Suppose we are given a commutative algebra A over a field F .

- Trivial d -algebra:
 - ▶ Suppose we are given a commutative algebra A over a field F .
 - ▶ Then we can turn A into a d -algebra by setting $d(a) = 0$ for all A .

- Trivial d -algebra:
 - ▶ Suppose we are given a commutative algebra A over a field F .
 - ▶ Then we can turn A into a d -algebra by setting $d(a) = 0$ for all A .
 - ▶ This is boring.

Examples of d -algebras

- Trivial d -algebra:
 - ▶ Suppose we are given a commutative algebra A over a field F .
 - ▶ Then we can turn A into a d -algebra by setting $d(a) = 0$ for all A .
 - ▶ This is boring.
- We are interested in noncommutative d -algebras.

Examples of d -algebras

- Trivial d -algebra:
 - ▶ Suppose we are given a commutative algebra A over a field F .
 - ▶ Then we can turn A into a d -algebra by setting $d(a) = 0$ for all A .
 - ▶ This is boring.
- We are interested in noncommutative d -algebras.

Quick sidenote: we also must assume that F is algebraically closed and $\text{char}(F) = 2$.

- Question: Can we find any noncommutative d -algebras?

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5: No
 - ▶ Dimension 6:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5: No
 - ▶ Dimension 6: No

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5: No
 - ▶ Dimension 6: No
 - ▶ Dimension 7:

Noncommutative d -Algebras

- Question: Can we find any noncommutative d -algebras?
- Solution Attempt: Try looking at d -algebras of small dimension.
 - ▶ Dimension 1: No
 - ▶ Dimension 2: No
 - ▶ Dimension 3: No
 - ▶ Dimension 4: No
 - ▶ Dimension 5: No
 - ▶ Dimension 6: No
 - ▶ Dimension 7: Yes!

7-dimensional d -Algebras

- We can actually construct a 7-dimensional d -Algebra.

7-dimensional d -Algebras

- We can actually construct a 7-dimensional d -Algebra.
- Our algebra A has basis (over F) $\{1, v_1, v_2, v_3, w_1, w_2, w_3\}$, and we define:

$$d(1) = d(v_1) = d(v_2) = d(v_3) = 0$$
$$d(w_1) = v_1 \quad \text{and} \quad d(w_2) = v_2 \quad \text{and} \quad d(w_3) = v_3$$

7-dimensional d -Algebras

- We can actually construct a 7-dimensional d -Algebra.
- Our algebra A has basis (over F) $\{1, v_1, v_2, v_3, w_1, w_2, w_3\}$, and we define:

$$d(1) = d(v_1) = d(v_2) = d(v_3) = 0$$
$$d(w_1) = v_1 \quad \text{and} \quad d(w_2) = v_2 \quad \text{and} \quad d(w_3) = v_3$$

- It has multiplication table:

\cdot	v_1	v_2	v_3	w_1	w_2	w_3
v_1	0	v_3	0	0	w_3	0
v_2	v_3	0	0	w_3	0	0
v_3	0	0	0	0	0	0
w_1	0	w_3	0	0	v_3	0
w_2	w_3	0	0	0	0	0
w_3	0	0	0	0	0	0

- A is noncommutative because $w_1 \cdot w_2 = v_3$, but $w_2 \cdot w_1 = 0$.

7-dimensional d -Algebras

- Question: Are there any other (up to isomorphism) 7-dimensional (noncommutative) d -algebras?

7-dimensional d -Algebras

- Question: Are there any other (up to isomorphism) 7-dimensional (noncommutative) d -algebras?
- For reference, there are infinitely many (up to isomorphism) commutative algebras of dimension 7 (Poonen, 2008).

7-dimensional d -Algebras

- Question: Are there any other (up to isomorphism) 7-dimensional (noncommutative) d -algebras?
- For reference, there are infinitely many (up to isomorphism) commutative algebras of dimension 7 (Poonen, 2008).
- Main Result:

Theorem

Up to isomorphism, there exists only one noncommutative d -algebra of dimension 7.

Acknowledgements

I would like to thank:

Acknowledgements

I would like to thank:

- My mentor Lucas Mason-Brown

Acknowledgements

I would like to thank:

- My mentor Lucas Mason-Brown
- Pavel Etingof

Acknowledgements

I would like to thank:

- My mentor Lucas Mason-Brown
- Pavel Etingof
- Siddharth Venkatesh

Acknowledgements

I would like to thank:

- My mentor Lucas Mason-Brown
- Pavel Etingof
- Siddharth Venkatesh
- The MIT PRIMES program and the MIT math department

Acknowledgements

I would like to thank:

- My mentor Lucas Mason-Brown
- Pavel Etingof
- Siddharth Venkatesh
- The MIT PRIMES program and the MIT math department
- My parents